3M ESPE TEMPORARY METAL CROWNS

Iso-Form[™] Crowns

Indications:

- · Short-term provisional coverage
- Premolars and molars

Tooth Preparation:

Prepare teeth using conventional techniques.

Crown Selection:

• Determine mesio-distal width. Select appropriate crown.

Trial fit for chamfer prep:

• Press gently to stretch crown cervix over finish line of preparation.

Trial fit for shoulder prep:

- Slightly flare the cervix of crown on the stretch block. Do not overexpand.
- Seat expanded crown over prep shoulder.











3M ESPE

Iso-Form[™] Crowns



Crown Preparation:

- If gingival blanching occurs during trial fit, trim crown using a 3M[™] ESPE[™] Crown Scissors.
- Reset in mouth and have patient establish occlusion.



- 99
- Use a sharp explorer to mark the marginal contour. Trim crown back to the line using a 3M[™] ESPE[™] Crown Scissors. Smooth and polish, if desired, using 3M[™] ESPE[™] Sof-Lex[™] Finishing and Polishing Discs and rubber wheel.
- Mix cement and fill crown, assuring voids are not present in cement.

Note: For custom margins and extended durability, line the crown with acrylic or resin material.



- **Crown Cementation:**
- Seat the crown and have the patient bite into occlusion.

2 of 3

3M ESPE TEMPORARY METAL CROWNS

Iso-Form[™] Crowns

- Burnish the cervical margin to the preparation before the cement sets completely.
- Remove excess cement with scaler or explorer. Floss interproximal areas.





3M ESPE Customer Hotline 1-800-634-2249 Please refer to instructions for more detailed information as well as precautionary and warranty information. 3M, ESPE, Iso-Form and Sof-Lex are trademarks of 3M or 3M ESPE AG. **3 of 3** 70-2009-2462-2 44-0019-9064-5 © 2003 3M